



Caramaschi, D., Hatcher, C., Mulder, R. H., Felix, J. F., Cecil, C. A. M., Relton, C. L., & Walton, E. (2020). Epigenome-wide association study of seizures in childhood and adolescence. *Clinical Epigenetics*, 12, [8 (2020)]. <https://doi.org/10.1186/s13148-019-0793-z>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1186/s13148-019-0793-z](https://doi.org/10.1186/s13148-019-0793-z)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via BMC at <https://clinicalepigeneticsjournal.biomedcentral.com/articles/10.1186/s13148-019-0793-z> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights


This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

RESEARCH

Open Access



Epigenome-wide association study of seizures in childhood and adolescence

Doretta Caramaschi^{1,2*} , Charlie Hatcher^{1,2}, Rosa H. Mulder^{3,4,5}, Janine F. Felix^{4,6,7}, Charlotte A. M. Cecil^{5,7}, Caroline L. Relton^{1,2} and Esther Walton^{1,2,8}

Abstract

The occurrence of seizures in childhood is often associated with neurodevelopmental impairments and school underachievement. Common genetic variants associated with epilepsy have been identified and epigenetic mechanisms have also been suggested to play a role. In this study, we analyzed the association of genome-wide blood DNA methylation with the occurrence of seizures in ~ 800 children from the Avon Longitudinal Study of Parents and Children, UK, at birth (cord blood), during childhood, and adolescence (peripheral blood). We also analyzed the association between the lifetime occurrence of any seizures before age 13 with blood DNA methylation levels. We sought replication of the findings in the Generation R Study and explored causality using Mendelian randomization, i.e., using genetic variants as proxies. The results showed five CpG sites which were associated cross-sectionally with seizures either in childhood or adolescence (1–5% absolute methylation difference at $p_{FDR} < 0.05$), although the evidence of replication in an independent study was weak. One of these sites was located in the *BDNF* gene, which is highly expressed in the brain, and showed high correspondence with brain methylation levels. The Mendelian randomization analyses suggested that seizures might be causal for changes in methylation rather than vice-versa. In conclusion, we show a suggestive link between seizures and blood DNA methylation while at the same time exploring the limitations of conducting such study.

Keywords: Seizures, Epilepsy, Epigenetics, DNA methylation, ALSPAC, Illumina 450 K, Mendelian randomization, Generation R Study

Background

Seizures are episodes of abnormal excessive or synchronous neuronal activity in the brain. When associated with a febrile illness, they affect 2–4% of children under 6 years of age in Europe and the USA, with the highest incidence in underdeveloped and rural areas reaching 14% in some areas [1, 2]. The most common age for seizures to occur is at 18 months of age and children that experienced seizures are at risk of developing epilepsy. The incidence of epilepsy in children ranges from 41 to 187/100,000 persons per year [2]. Seizures and epilepsy are associated with neurodevelopmental conditions, such as autism spectrum disorders [3], attention-deficit hyperactivity disorder, and cognitive

impairment [4–7]. Moreover, epilepsy with or without intellectual impairment is associated with low academic achievement [8].

The two largest genome-wide association studies to date (~ 8600 individuals with epilepsy versus ~ 26000 controls and ~ 15,200 individuals with epilepsy versus ~ 29,600 controls) have identified a total of 24 genetic variants associated with epilepsy [9, 10]. Some of these loci are located in the proximity of candidate genes for epilepsy, for instance those coding for ion-channel subunits, and their relevance for epilepsy is supported by other research in humans and animals. As it is likely that other factors may also underlie the disease, it has been suggested that epigenetic mechanisms such as DNA methylation are also involved in the onset of seizures [11]. In line with this hypothesis, genetic markers for epilepsy were found to be enriched in histone modification markers, suggesting epigenetic regulation of gene transcription [9]. The association between seizures and DNA

* Correspondence: D.Caramaschi@bristol.ac.uk

¹Bristol Medical School, Population Health Sciences, University of Bristol, 5 Tyndall Avenue, Bristol BS8 1UD, UK

²Medical Research Council Integrative Epidemiology Unit, University of Bristol, Oakfield House, Oakfield Grove, Bristol BS8 2BN, UK

Full list of author information is available at the end of the article



methylation has been investigated in studies involving humans and other animals, although these studies relied on small sample sizes or on a candidate gene approach. There are different types of epilepsy syndromes depending on the age of onset (e.g., childhood or adolescence), whether the seizures are predominantly characterized by a focal or generalized onset, and whether there are known causes (e.g., genetic or trauma) [12]. Mesial temporal lobe epilepsy is one of the most common and most studied forms of epilepsy [13]. A recent study comparing blood DNA methylation in 30 adult patients with mesial temporal lobe epilepsy and 30 controls identified 216 differentially methylated sites between the two groups, including sites on genes involved in ion binding and metabolic activity [14]. DNA methylation differences have also been observed in lymphoblastoid cell lines derived from epileptic patients, both when DNA methylation was measured globally using antibody capture and in the *BRD2* gene promoter [15]. Another study that reanalyzed these data discovered differential DNA methylation in non-coding RNAs [16]. Furthermore, alterations in DNA methylation were present in the hippocampus of epileptic patients compared to controls [17]. A study adopting a rat model of chronic epilepsy corroborated these findings by revealing genome-wide differences in DNA methylation compared to control rats [18].

Typically, in association studies, it is difficult to assess the causality of any identified association due to the potential for confounding and/or for reverse causation. Socioeconomic status, for instance, is associated with genome-wide changes in DNA methylation [19] and with increased risk for seizures/epilepsy [20], suggesting that socioeconomic factors could be confounding the association between DNA methylation and seizures. With respect to reverse causation, case-control studies that examined DNA methylation after epilepsy had already been diagnosed might have observed changes that were directly caused by the seizure events. For instance, laboratory animal studies have shown altered gene expression after induced seizures [21]. Mendelian randomization, a technique that uses the genetic information associated with an exposure to estimate the causal effect of the exposure on an outcome, can circumvent these limitations under certain assumptions [22].

In this study, we (1) investigated the genome-wide association of DNA methylation with the occurrence of seizures from birth throughout childhood and adolescence in peripheral blood samples from a prospective birth cohort, (2) carried out replication analyses in an independent study sample, (3) explored the correspondence with brain tissue DNA methylation at the same genomic locations to investigate whether these associations have neurodevelopmental relevance, and (4)

performed bi-directional Mendelian randomization to unravel causal links between peripheral blood DNA methylation and seizures. Finally, we explored the potential health consequences of a seizure-associated DNA methylation profile. For an overview on our analysis plan, see Fig. 1.

Materials and methods

Study population

The discovery analyses were conducted in the Avon Longitudinal Study of Parents and Children (ALSPAC), a large prospective cohort study that recruited 14,541 pregnant women, resident in Avon, UK with expected delivery dates between the 1st of April 1991 and the 31st of December 1992 [23, 24]. Of these initial pregnancies, there were 14,062 live births and 13,988 children who were alive at 1 year of age. The study website contains details of all the data that are available through a fully searchable data dictionary (<http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary/>).

Written informed consent has been obtained for all ALSPAC participants. Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees.

Seizure data

The diagnosis of seizures and epilepsy is usually based on the pattern of seizures, age of onset, and electroencephalographic and imaging features, and potentially the history of comorbidities as well as genetic and/or metabolic screening [25]. In our ALSPAC population study, we relied on data taken from questionnaires administered to the mothers at four time points: 18 months, 8 years, 11 years, and 13 years. The following questions were asked at each of these ages:

18 months: "Has he/she ever had any form of convulsion/fit/seizure or other turn in which consciousness was lost or any other part of the body made an abnormal movement?",

8 years: "Has child had a convulsion/fit/seizure since 7th birthday?",

11 years: "Has child had a convulsion/fit/seizure where consciousness or abnormal movement was lost since 9th birthday?", and

13 years: "Has she ever had a seizure, fit or a convulsion?".

This allowed us to address proximal and distal (i.e., lifetime until age 13) associations between DNA methylation and experience of seizures. The seizures reported in our study could have included any type of seizure that might not be considered epilepsy, such as febrile seizures typically experienced before 6 years of age and single occurrences provoked by trauma. Alternatively, they could be recurring seizures of the types typical of an epilepsy




Fig. 1 Analysis overview

syndrome. However, we did not have access to the information on the clinical diagnoses in relation to epilepsy in the study participants.

DNA methylation data

In ALSPAC, blood from 1018 mother-child pairs were selected for analysis as part of the Accessible Resource for Integrative Epigenomic Studies (ARIES, <http://www.ariesepigenomics.org.uk/>) [26]. Following DNA extraction, samples were bisulphite converted using the Zymo EZ DNA Methylation™ kit (Zymo, Irvine, CA, USA), and genome-wide methylation was measured using the Illumina Infinium HumanMethylation450 (HM450) BeadChip. The

arrays were scanned using an Illumina iScan, with initial quality review using GenomeStudio. ARIES was pre-processed and normalized using the *meffil* R package [27]. ARIES consists of from mother-child pairs measured at five time points (three time points for children: birth, childhood, and adolescence; and two for mothers: during pregnancy and at middle age), although only children's profiles were used in the current study. Low quality profiles were removed from further processing, and the remaining 4593 profiles were normalized using the Functional Normalization algorithm [28] with the top 10 control probe principal components. Full details of the pre-processing and normalization of ARIES have been described previously [27]. Further pre-

processing specific to the current study included removal of probes not passing background detection ($p > 0.05$) and probes on the X or Y chromosome. To reduce the impact of outliers, we set methylation data points outside the 3× inter-quartile-range from the 25th and the 75th percentiles to missing. The total number of probes available for analyses were $N = 468,828$ at birth; $N = 471,092$ at childhood; and $N = 470,480$ at adolescence.

Epigenome-wide association analyses (EWAS)

In ALSPAC, the final sample size at birth was $N = 822$ (25 cases and 797 controls); at childhood $N = 848$ (7 cases and 841 controls) and $N = 813$ (12 cases and 801 controls) at adolescence. The final sample only contained singletons and no siblings. Only 2–3 cases overlapped across time points (Additional file 1: Figure S1). This is to be expected as certain types of seizures, such as febrile ones, are more common in the first years of life and children diagnosed with epilepsy might have received treatment to prevent further seizures. Moreover, the age of onset of some forms of epilepsy is between childhood and adolescence [29], explaining why some children might have had their first seizures in their adolescence. Fifty-eight out of a total of $N = 817$ adolescents reported a lifetime experience of seizures.

We carried out four analyses. In analysis 1–3, we modelled methylation at birth (model 1), childhood (model 2), and adolescence (model 3) as the outcome and seizure status (measured closest to each methylation time point) as the exposure. To investigate the temporal sensitivity of associations, we ran a final analysis (model 4), in which we modelled lifetime seizure status (ever/never) at 13 years as the exposure and methylation at adolescence as the outcome (model 4). In all models (including birth), methylation was defined as the outcome regardless of temporal order to keep model estimates consistent and comparable. These EWAS were carried out in R version 3.3.1 using the CpGassoc package [30].

All models were adjusted for age (when DNA methylation samples were taken), sex, prenatal maternal smoking (Yes/No), and maternal education (university degree Yes/No), each derived from ALSPAC maternal and childhood questionnaires. The model using cord blood data was additionally adjusted for gestational age and birthweight. Unknown confounders and batch were adjusted using surrogate variable (SV) analysis [31]. Additionally, we adjusted for cell counts using the Houseman method for the childhood and adolescence timepoints [32] and the Andrews and Bakulski method for cord blood [33].

To summarize, the following models were applied:

Model 1: Methylation (cord) ~ seizure status (18 months) + age + sex + birthweight + gestational age + maternal prenatal smoking + maternal education +

nucleated red blood cells + granulocytes + monocytes + natural killer cells + B cells + CD4(+)T cells + CD8(+)T cells + SV1 + ... + SV15

Model 2: Methylation (childhood) ~ seizure status (8 years) + age + sex + maternal prenatal smoking + maternal education + granulocytes + monocytes + natural killer cells + B cells + CD4(+)T cells + CD8(+)T cells + SV1 + ... + SV13

Model 3: Methylation (adolescence) ~ seizure status (11 years) + age + sex + maternal prenatal smoking + maternal education + granulocytes + monocytes + natural killer cells + B cells + CD4(+)T cells + CD8(+)T cells + SV1 + ... + SV14

Model 4: Methylation (adolescence) ~ seizure status (ever/never at 13 years) + age + sex + maternal prenatal smoking + maternal education + granulocytes + monocytes + natural killer cells + B cells + CD4(+)T cells + CD8(+)T cells + SV1 + ... + SV14

To correctly multiple testing, we present both Bonferroni (0.05/number of probes) and FDR-corrected results.

Replication analyses

All CpG sites that were associated with seizures below at least an FDR-correction threshold in ALSPAC were analyzed in an independent cohort to assess replication. The Generation R Study is a population-based prospective cohort study conducted in Rotterdam, the Netherlands, that recruited 9778 pregnant women with an expected delivery date between April 2002 and January 2006. A total of 9749 children were born from these pregnancies, and extensive data and biological samples are available from the children and their mothers [34]. DNA methylation was measured in peripheral blood of 469 children aged 6 years (all singletons), using the Infinium HumanMethylation450 (HM450) BeadChip as in ALSPAC. Preparation and normalization of the BeadChip array was performed according to the CPACOR workflow in R [35] and methylation data points lower than the 25th percentile – 3 × IQR and higher than the 75th percentile + 3 × IQR were excluded. In Generation R, seizure events were measured using the answer “yes” to the question “During the past 5/6 years, did your child ever have a seizure/febrile convulsion?” asked to the mothers when the children were 6 years of age. The final sample size was $N = 432$, with 19 participants affected by seizures and 413 unaffected (Additional file 1: Table S1). Linear models similar to model 2 were run in Generation R on EWAS FDR-corrected methylation sites and on *BDNF* probes. The covariates were measured and categorized similarly to the analyses carried out in ALSPAC. The results in ALSPAC and Generation R were also meta-analyzed using METAL [36], using inverse variance weighting.

Mendelian randomization analyses

To assess the causal link between DNA methylation and the occurrence of seizures, we performed two-sample Mendelian randomization (MR) using the EWAS results for the CpG sites with FDR-corrected p values < 0.05 in ALSPAC. Two-sample MR was performed using the MR-Base online platform (<http://www.mrbase.org/>, last accessed 06-06-2018) [37], the MRInstruments R package (<https://github.com/MRCIEU/MRInstruments>, last accessed 06-06-2018), and the TwoSampleMR R package (<https://github.com/MRCIEU/TwoSampleMR>, last accessed 06-06-2018).

Three MR analyses were performed. (1) To investigate the causal effect of DNA methylation on the risk of epilepsy, we performed two-sample MR with DNA methylation as exposure and a diagnosis of epilepsy as outcome. For the genotype-exposure associations, we searched for methylation quantitative trait loci (mQTLs), i.e., genetic variants that are associated in *cis* with DNA methylation (i.e., within 1 Mb either side from the CpG site), using the mQTL database (<http://www.mqtl.org/>, last accessed 06 June 2018) [38], restricting the search to the time point, at which the CpG site was associated with seizure status. For the genotype-epilepsy association, we used the summary statistics for ICD-9 and ICD-10 codes for epilepsy or seizure in MR-Base generated on UKBiobank data (last accessed 04 December 2018). (2) To analyze the causal effect of risk for seizure/epilepsy on DNA methylation (i.e., reverse causation), we performed two-sample MR with epilepsy diagnosis or febrile/vaccine-related seizures as exposure and DNA methylation as outcome. For the genotype-exposure associations, we used the summary statistics for the genome-wide significant SNPs from a published GWAS meta-analysis on all epilepsies, focal epilepsy, and genetic generalized epilepsy [10] and from a published GWAS on febrile and MMR-vaccine-related seizures (focussing on 6 replicated genome-wide significant SNPs in Table 1 of the original article) [39]. Odds ratios (OR) and confidence intervals were reverted to log-odds and standard errors to use in the two-sample MR analysis. The summary statistics for the genotype-outcome associations were drawn from the mQTL database (<http://www.mqtl.org/>, last accessed 06 June 2018) [38]. (3) To analyze the causal effect of seizure-susceptible DNA methylation (exposure) on other health outcomes, we first performed a hypothesis-free PheWAS using the mQTLs for CpG sites associated with seizures and the MR-Base online PheWAS tool to screen for potentially affected health outcomes. Then, we performed two-sample MR with methylation as exposure and the health outcomes identified by the PheWAS to estimate the magnitude of the effect. We also performed a hypothesis-driven two-sample MR on other

neurodevelopmental outcomes previously found to associate with seizures, i.e., autism, intelligence, and education.

Results

For a flowchart and overview of all results, see Additional file 1: Figure S2.

Sample description

At 18 months, there were $n = 25$ children with the experience of seizures since birth and $n = 797$ without (Table 1). Groups were comparable with respect to sex, birthweight, gestational age, as well as maternal education and smoking behavior during pregnancy. During childhood, $n = 7$ children had experienced seizures between 7 and 8 years of age while 841 had not. Age at blood draw was slightly higher in the children with seizures (Table 1). Approaching adolescence, seizures had been reported in $n = 12$ children, while $n = 801$ children had not experienced seizures between 9 and 11 years of age. There were slightly more females in the seizure group. Fifty-eight adolescents reported a lifetime experience of seizures, while $n = 759$ had never experienced seizures. Groups were comparable with respect to maternal smoking behavior, maternal education, and age at blood draw. For correlation plots on seizure rates and all covariates included in the final models specific to each time point, see Additional file 1: Figures S3

Epigenome-wide association analyses

We did not identify any CpG sites that fell below a Bonferroni or FDR-adjusted P value threshold at birth (Table 2 and Additional file 1: Figure S4). In childhood, two CpG sites were associated with seizure status at an FDR-corrected p value < 0.05 (cg10541930: beta = -0.010 , SE = 0.002, uncorrected p value = 4.32×10^{-8} and FDR = 0.020; cg25557432: beta = 0.014, SE = 0.003, uncorrected p value = 1.82×10^{-7} and FDR = 0.043), of which cg10541930 was also below the Bonferroni threshold ($0.05/470,489 = 1.06 \times 10^{-7}$). The first CpG is located in an intergenic region, at the transcription start site for a non-coding RNA, while the second lies upstream of the gene *MACROD2* involved in DNA repair.

In adolescence, we found one CpG site that fell below a Bonferroni threshold of $0.05/470,479 = 1.06 \times 10^{-7}$, cg13974632 (beta = 0.053, SE = 0.010, uncorrected p value = 5.55×10^{-8} and FDR = 0.026), while another two CpG sites passed an FDR threshold only (cg15810326: beta = 0.014, SE = 0.003, uncorrected p value = 1.19×10^{-7} and FDR = 0.028; cg16983916: beta = -0.056 , SE = 0.011, uncorrected p value = 1.79×10^{-7} and FDR = 0.028; Fig. 2a). The first CpG site is located in the first exon of the brain-derived neurotrophic factor (BDNF). The experience of seizures was associated with increased DNA methylation at this site (Fig. 2b). The

Table 1 ARIES sample characteristics

Characteristic	Birth			Childhood			Adolescence			Adolescence (lifetime exposure)		
	With seizures	Without seizures	Total	With seizures	Without seizures	Total	With seizures	Without seizures	Total	With seizures (ever)	Without seizures (ever)	Total
N	25	797	822	7	841	848	12	801	813	58	759	817
Sex (%F)	44%	51%	51%	ND	50%	51%	ND	51%	52%	52%	51%	51%
Age at methylation in years (SD)	NA			7.69 (0.36)	7.44 (0.13)	7.45 (0.14)	17.65 (0.86)	17.11 (1.04)	17.12 (1.04)	17.16 (1.01)	17.10 (1.05)	17.10 (1.04)
Age at seizure assessment in years (SD)	1.54 (0.12)	1.52 (0.06)	1.52 (0.07)	8.60 (0.03)	8.66 (0.16)	8.66 (0.16)	11.70 (0.08)	11.71 (0.12)	11.71 (0.12)	13.12 (0.09)	13.15 (0.17)	13.43 (1.05)
Birthweight in g (SD)	3498.80 (545.54)	3495.08 (478.18)	3495.20 (480.00)	NA			NA			NA		
Gestational age in weeks (SD)	39.80 (1.50)	39.55 (1.48)	39.56 (1.48)									
Maternal smoking (%Y)	ND	13%	14%	ND	12%	13%	ND	13%	13%	14%	13%	13%
Maternal education (%Uni)	32%	20%	21%	ND	21%	21%	ND	22%	22%	28%	22%	22%

ND not disclosed due to cohort restrictions, NA not applicable

Table 2 EWAS results. Top CpG sites for the three time-points: birth, childhood, and adolescence (cross-sectional and lifetime exposure)

Time point	ProbeID	Beta	SE	P value	FDR	Bonferroni	Chromosome	Position	Gene
Birth	cg07504545	− 0.042	0.008	2.32×10^{-07}	0.109	0.109	1	203456019	PRELP
	cg04977770	− 0.054	0.011	7.15×10^{-07}	0.168	0.335	17	79846763	THOC4
	cg23625106	− 0.023	0.005	1.47×10^{-06}	0.193	0.689	8	61789727	
	cg08461451	− 0.036	0.008	1.73×10^{-06}	0.193	0.812	19	2295092	LINGO3
	cg06367149	0.057	0.012	2.08×10^{-06}	0.193	0.973	15	61254575	RORA
	cg13396019	− 0.026	0.005	2.57×10^{-06}	0.193	1.000	1	220564510	
Childhood	cg10541930	− 0.010	0.002	4.32×10^{-08}	0.020	0.020	10	131909085	
	cg25557432	0.014	0.003	1.82×10^{-07}	0.043	0.086	20	13976117	MACROD2
	cg11317171	0.002	0.000	1.34×10^{-06}	0.190	0.633	19	5623025	SAFB2;SAFB
	cg06447795	0.090	0.019	1.62×10^{-06}	0.190	0.761	20	36793996	TGM2
	cg02389501	0.078	0.016	2.59×10^{-06}	0.191	1.000	7	93757948	
	cg13647052	0.018	0.004	2.82×10^{-06}	0.191	1.000	12	2800382	CACNA1C
Adolescence	cg13974632	0.053	0.010	5.55×10^{-08}	0.026	0.026	11	27740813	BDNF
	cg15810326	0.014	0.003	1.19×10^{-07}	0.028	0.056	4	148605127	PRMT10
	cg16983916	− 0.056	0.011	1.79×10^{-07}	0.028	0.084	7	156159713	
	cg11510269	0.007	0.001	5.18×10^{-07}	0.050	0.244	12	123011761	RSRC2;KNTC1
	cg17130518	0.052	0.010	5.33×10^{-07}	0.050	0.251	7	76825482	CCDC146;FGL2
	cg06222062	0.095	0.019	6.79×10^{-07}	0.053	0.319	1	20396690	PLA2G5
Lifetime	cg20162381	0.004	0.001	9.22×10^{-07}	0.159	0.434	12	3310097	TSPAN9
	cg20156774	− 0.028	0.006	1.52×10^{-06}	0.159	0.717	6	32803500	TAP2
	cg12888660	0.003	0.001	1.72×10^{-06}	0.159	0.811	12	122018449	KDM2B
	cg15031661	0.006	0.001	1.91×10^{-06}	0.159	0.897	1	240256603	FMN2
	cg17233506	− 0.031	0.006	2.21×10^{-06}	0.159	1.000	17	46608293	HOXB1
	cg18108087	0.010	0.002	2.32×10^{-06}	0.159	1.000	1	59762112	FGGY




Fig. 2 **a** Miami plot displaying the EWAS results by chromosome in adolescence. Positive values on y-axis indicate $-\log(p\text{ values})$ of hypermethylated sites, whereas negative values on the y-axis indicate $-\log(p\text{ values})$ of hypomethylated sites (the sign of the y-axis values has been changed to reflect this). Bonferroni cut-off line in red. **b** Boxplot of methylation levels at the *BDNF*-linked CpG cg13974632 (adjusted for covariates). **c** Causal estimate for the effect of genetic generalized epilepsy on cg13974632 (*BDNF*). Individual SNP results in black and overall causal estimates in red. **d** Leave-one-out analysis of the causal estimate for the effect of genetic generalized epilepsy on cg13974632 (*BDNF*). IVW inverse variance weighted

other two CpG sites were located in the first exon of Protein Arginine Methyltransferase 10 (*PRMT10*) and in an intergenic region, respectively.

A sensitivity analysis where we adjusted for fewer cell types to avoid overfitting (i.e., we omitted CD8+ T cell proportions) showed similar association estimates in the probes at $FDR < 0.05$ in the main EWAS (Additional file 1: Table S2).

No CpG site could be identified to be associated with lifetime seizure experience. Inspection of QQ plots and lambdas close to 1 gave little indication for an inflation of test statistics (Additional file 1: Figure S5).

When testing the association of the significant CpGs at childhood or adolescence across the other age ranges, the effect sizes were much smaller, not significant after correction for multiple comparisons, and for some CpGs in the opposite direction. (Additional file 1: Table S3).


Replication in the Generation R Study

For replication, we focused on five CpG sites: two that passed FDR correction in childhood and three that passed FDR correction in adolescence. As one of the

CpG sites in adolescence was located in the gene *BDNF*, which is a key driver in neuronal growth and has been repeatedly linked to epilepsy [40, 41], we expanded our search space to include all CpG sites annotated to *BDNF* ($n = 73$).

None of the five CpG sites were associated with seizures in Generation R during childhood. Although the direction of effect suggested some degree of concordance, all but one beta coefficient were a factor of 10 smaller and p values ranged between 0.1 and 0.8. (Additional file 1: Table S4 and Fig. 3). When ALSPAC and Generation R results for these five probes were meta-analyzed together, all CpGs except for cg16983916 showed evidence of methylation differences ($p\text{ value} < 0.05/77 = 0.0006$, Additional file 1: Table S4).

Investigating all 73 CpG sites annotated to *BDNF*, none replicated based on a correction for 73 tests. We observed only a weak correlation of all 73 regression betas between cohorts ($\rho = 0.046$, $p\text{ value} = 0.70$ based on adolescence results in ALSPAC and childhood results in Generation R; Additional file 1: Table S4 and Additional file 1: Figure S6). However, five CpG sites



were significant at a nominal level. When ALSPAC and Generation R results for these 73 probes were meta-analyzed together, there was evidence for methylation differences in 2 CpGs ($p < 0.05/77 = 0.0006$, Additional file 1: Table S4). These were *cg13974632*, the top hit from the EWAS, and *cg15313332*, 20Kb upstream.

Cross-tissue concordance in DNA methylation

We queried three independent databases to investigate blood-brain concordance in DNA methylation for all five CpG sites that passed FDR correction in ALSPAC. Based on data from more than 122 pre-mortem blood samples and paired post-mortem brain tissue [42], cross-tissue correlation was strongest for *BDNF*_{*cg13974632*} ($r = 0.39$) between blood and brain tissue from the entorhinal cortex, followed by tissue from the prefrontal cortex ($r = 0.27$); Additional file 1: Figure S7A-E. Compared to *BDNF*_{*cg13974632*}, the remaining four CpG sites showed correlations, which were generally weaker for tissue from the prefrontal and entorhinal cortex. Cross-tissue correlations based on a smaller sample of 16 individuals [43] reported different correlation profiles (Additional file 1: Figure S7F-G). In this dataset, the blood-brain correlation was strongest for *MACROD2*_{*cg25557432*} in Brodmann Area BA20 (temporal cortex, $\rho = 0.48$) and BA7 (parietal cortex, $\rho = 0.43$), and for *cg15810326* in BA20 ($\rho = 0.31$), whereas there was little evidence of a

positive correlation for the other sites in either BA10 (prefrontal cortex), BA20 or BA7.

Cross-tissue blood-brain correlations in a third dataset of 12 epilepsy patients [44] were consistent with the larger dataset, although not discriminating between brain regions (*BDNF*_{*cg13974632*}: $\rho = 0.28$, $p = 0.42$; *PRMT10*_{*cg15810326*}: $\rho = -0.0$, $p = 0.94$; *cg16983916*: $\rho = 0.09$, $p = 0.81$; *cg10541930*: $\rho = 0.32$, $p = 0.36$; *MACROD2*_{*cg25557432*}: $\rho = 0.45$, $p = 0.19$). Although the correlation coefficients were similar to the larger dataset ($N = 122$), in this smaller dataset ($N = 12$), there was only 9% power to detect a correlation as low as 0.2 at $\alpha = 0.05$ and 26% power to detect a correlation of 0.4.

Based on data available via the Genotype-Tissue Expression (GTEx; www.gtexportal.org) project, we investigated tissue-specific gene expression for genes linked to the five FDR-corrected CpG sites. *BDNF* appeared to be expressed in the brain and other tissues with highest expression in the cerebellum, while very low expression was found in blood. *MACROD2* is predominately expressed in lymphocytes; *PRMT10* is mainly expressed in the ovaries (Additional file 1: Figure S8).

Mendelian randomization analyses

Uni- and bi-directional two-sample MR was performed to investigate the effect of DNA methylation on seizure occurrence and the effect of seizures on DNA methylation. We used genetic associations with febrile and vaccine-related seizures and genetic associations with epilepsy. The latter more generally includes the types of seizures observed in our study (e.g., febrile and non-febrile) [45]. As full genome-wide summary statistics were only available for epilepsy, but not for febrile and vaccine-related seizures, we could not perform the two-sample MR to estimate the causal effects of methylation on seizures. We identified only one *cis*-mQTL, which could be used as instrument for DNA methylation. In detail, the SNP rs10258194 was associated in *cis* with *cg16983916* (effect allele = T, $\beta = 0.25$, $SD = 0.04$, $p = 2.33 \times 10^{-10}$) after excluding other SNPs due to linkage disequilibrium. For the other CpG sites, either *trans*-associations (further than 1 Mb from the CpG site) or no associations were identified. Two-sample MR indicated only weak evidence for causal effects of DNA methylation at *cg1698369* (Additional file 1: Table S5) on epilepsy.

For the reverse (i.e., epilepsy/seizure affecting DNA methylation), there were 9 SNPs to be used as instruments for epilepsy from a previous GWAS meta-analysis on all epilepsies, focal epilepsy, and genetic generalized epilepsy, although only up to four SNPs were used in any one analysis due to availability of summary statistics. Six SNPs were identified as instruments for febrile/vaccine-related seizures, although only 5 were used.

Additional file 1: Table S6 shows the results of the two-sample MR analysis performed using different methods to investigate the causal effect of epilepsy on DNA methylation at the five CpG sites identified in the EWAS. For cg13974632 (*BDNF*), there was some evidence for a positive association of genetic generalized epilepsy with increased DNA methylation using the weighted median method (Fig. 2c, d). All methods, including MR-Egger, suggested a positive effect of genetic generalized epilepsy on cg13974632, although these analyses were based on only 3 genetic instruments, and confidence intervals were large, particularly for MR-Egger. The leave-one-out analysis indicated that this effect was not driven by a particular genetic variant, providing little evidence for a violation of MR assumptions. This association did not survive a correction for multiple testing and seemed to be specific for genetic generalized epilepsy (i.e., the effect was not reproduced using focal epilepsy or “any epilepsy” as an exposure). There was no evidence for associations with any other CpG sites.

The two-sample MR analysis of the effects of febrile and vaccine-related seizures on methylation did not show enough evidence of a causal association (Additional file 1: Table S7).

To test the effect of seizure-associated methylation on other health outcomes, we scanned for potentially relevant health traits by performing a PheWAS (association of genotype with all available outcomes) and, subsequently, two-sample MR using the only mQTL available, rs10258194. As we used only one instrument, we could not differentiate whether the associations were due to causal effects or horizontal pleiotropy (i.e., the genetic variant has an effect on the health outcome outside of its effect on DNA methylation at the specific CpG). The analysis revealed little evidence for associations between rs10258194 and other health outcomes, both when analyzed in a hypothesis-free screening PheWAS across all available outcomes (Additional file 1: Table S8) and

when analyzed in a two-sample MR with specified outcomes (top outcomes from PheWAS and neurodevelopmental outcomes, Table 3).

Discussion

In this study, we observed associations between blood DNA methylation and the occurrence of seizures in a longitudinal pregnancy cohort study based in the UK. Effects were specific to childhood and adolescence, with little evidence for a relationship at birth or for lifetime exposure to seizures. However, associations did not replicate in an independent study sample based in the Netherlands. The results are summarized in Additional file 1: Figure S2.

This study has a number of strengths. First, in both ALSPAC and Generation R cohorts, the information on seizures was provided by the parents near the time of occurrence, therefore reducing measurement error and the possibility of recall bias. Secondly, repeated blood sampling at different ages in ALSPAC, including birth, allowed age-specific cross-sectional analyses. Thirdly, these studies have collected extensive information from obstetric records and reported socioeconomic factors allowing adjustment for potential confounders, including birth weight as well as maternal smoking during pregnancy and maternal education. Finally, we used a Mendelian randomization approach as an alternative method to control for unmeasured confounding and examine the direction of observed associations.

Our results in the discovery cohort suggest a link between *BDNF*, a neurotrophin that is highly expressed in the brain, and seizures and epilepsy, at a site where blood and brain DNA methylation levels show correspondence. Mendelian randomization analyses suggest a potential causal effect of seizures on DNA methylation in the *BDNF* gene. Although the implications of the association with *BDNF* are interesting, the association was not replicated. Studies conducted in animal models of

Table 3 Mendelian randomization analysis. Effects of DNA methylation at cg16983916 on non-epilepsy outcomes (PheWAS and candidate outcomes)

Outcome	Reference	N	Beta ^a	S.E.	P value (uncorrected)
Weight	UK Biobank ^b	336227	0.016	0.009	0.067
Trunk fat-free mass	UK Biobank ^b	331030	0.025	0.006	3.73×10^{-05}
Standing height	UK Biobank ^b	336474	0.029	0.007	2.90×10^{-05}
Years of schooling	Okbay et al. [46]	182286	0.008	0.008	0.317
Childhood intelligence	Benyamin et al. [47]	12441	0.013	0.051	0.797
Autism	Smoller et al. [48]	29415	− 0.008	0.113	0.946
Fluid intelligence score	UK Biobank ^b	108818	0.119	0.036	0.001
College or University degree	UK Biobank ^b	334070	0.013	0.004	0.004

^aWald ratio

^bUnpublished summary statistics from the UK Biobank. https://github.com/Nealelab/UK_Biobank_GWAS and <http://www.nealelab.is/blog/2017/9/11/details-and-considerations-of-the-uk-biobank-gwas>

epilepsy (reviewed in [49]) observed an upregulation of *BDNF* immediately after experimentally induced seizures. A study conducted on hippocampal tissue from 40 adult patients affected by mesial-temporal lobe epilepsy showed increased or decreased *BDNF* expression, compared to healthy individuals, depending on the region investigated and on the presence of psychiatric comorbidities [50]. Similarly, four isoforms of *BDNF* were found to be highly expressed in brain hippocampal tissue from adult epileptic patients compared to healthy controls, although the effect was not explained by changes in DNA methylation measured in the promoters of isoforms IV and VI [41]. It is to be noted that the association reported in the current study was located further upstream within the first intron of isoforms I, II, and III, based on the latest gene characterization [41, 51]. Moreover, a recent family study investigating genome-wide DNA methylation in peripheral blood, based on 15 trios of parents and their offspring, where the child and one parent, but not the other, were affected by generalized genetic epilepsy, found evidence of neurotrophins involvement, particularly *BDNF*, which was both hyper- and hypomethylated [52]. In our study, we observed hypermethylation in the *BDNF* gene (in the promoter or within introns, depending on the isoform), which would suggest decreased expression. This is in apparent contrast with some of the previous studies, but in line with our Mendelian randomization analysis that also showed some evidence of seizure-induced hypermethylation in the *BDNF* gene when using generalized genetic epilepsy as the exposure. The apparently contrasting findings in the *BDNF* gene and the lack or reproducibility of this association could be explained by the fact that *BDNF* expression in blood is very low, but follow-up investigations in biological systems with multi-tissue characterization are required to elucidate this further.

We also show potential associations with other methylation sites that have not been previously observed, specifically in the *MACROD2*, the *PRMT10* genes, and two intergenic sites. Single nucleotide polymorphisms within *MACROD2* have previously been associated with autism, although with rather weak evidence [53], and other brain-related traits such as intelligence and mathematical abilities [54]. Individuals with *de novo* mutations in *FBXO11*, an analogous of *PRMT10*, have been reported to show intellectual disability and autism [55]. Although both genes are predominantly expressed in non-neural tissues, these studies suggest that *MACROD2* and *PRMT10* could play a role in brain functioning, and their methylation status could plausibly be involved in seizures. However, as these associations were not replicated, the involvement of DNA methylation in blood at these sites needs to be further investigated.

The results of this study have to be seen in light of the following limitations. First, despite this being the largest epigenome-wide association study in childhood/adolescence, the sample size was small considering the low prevalence of seizures in the general population. Second, our initial findings did not replicate in data from an independent cohort. However, this could also indicate that the associations are specific to the time window examined in the discovery sample. For instance, the association with *BDNF* methylation was specific to the adolescence time point, while in Generation R data was limited to childhood. Thirdly, we have relied on DNA methylation measured in blood whereas seizures occur within the brain. Availability of brain tissue for epidemiological research is very limited, and future studies could follow-up our findings in animal models or post-mortem brain tissues. Finally, the Mendelian randomization analyses relied on a small number of mQTLs (3 for epilepsy to *BDNF* methylation and 1 for methylation to epilepsy and other health outcomes) and therefore did not allow for sensitivity analyses aimed at ruling out bias due to horizontal pleiotropy.

Conclusions

In conclusion, we show that the evidence of a link between seizures and blood DNA methylation in childhood and adolescence is weak. Our study highlights the challenges of conducting epigenome-wide association studies of seizures across different developmental periods and warrants a careful analysis of the data in view of the limitations of such study.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s13148-019-0793-z>.

Additional file 1: Figure S1. Overlap of cases across time points. **Figure S2.** Overview of results. All models were adjusted for covariates as specified in the main text (including cell composition and surrogate variables). In MR analyses, epilepsy was chosen as a trait due to the availability of GWAS summary data as well as febrile and MMR-vaccine-related seizures. **Figure S3.** Plots displaying bivariate correlations between all variables included in the final models: (A) at birth, (B) during childhood and (C-D) adolescence (cross-sectional and lifetime seizure exposure, respectively). **Figure S4.** Miami plots displaying EWAS results: (A) at birth, (B) during childhood and (C-D) adolescence (cross-sectional and lifetime seizure exposure, respectively). The sign of the y-axis (-log(P-values)) has been changed to indicate positive and negative changes in DNA methylation. Bonferroni cut-off line in red. **Figure S5.** Quantile-quantile plots displaying potential test statistic inflation in the final models (A) at birth, (B) during childhood and (C-D) adolescence (cross-sectional and lifetime seizure exposure, respectively). Lambda was calculated using the regression method. **Figure S6.** Differences in DNA methylation according to experienced seizures in ALSPAC and Generation R (top panel) in the *BDNF* gene. The bottom panel shows the location of the transcripts reverse strand. **Figure S7.** Cross-tissue correspondence of CpG sites, passing FDR correction in ALSPAC, based on data available at A-E) <https://epigenetics.essex.ac.uk/bloodbrain> and F-G) <https://redgar598.shinyapps.io/BECon>. PFC = prefrontal cortex; STG = superior temporal

gyrus; EC = entorhinal cortex; CER = cerebellum. **Figure S8.** Tissue-specific expression of BDNF, MACROD2 and PRMT10, based on data available at www.gtportal.org. **Table S1.** Sample descriptives of Generation R. **Table S2.** Association estimates for FDR<0.05 probes re-analyzed with adjustment for 5 cell types. **Table S3.** Association of DNA methylation and the occurrence of seizures at CpGs with FDR<0.05 across all models. **Table S4.** Replication in Generation R and meta-analysis. **Table S5.** Two-sample MR analysis of the effect of DNA methylation on epilepsy (method=Wald ratio). **Table S6.** Two-sample MR analysis of the effect of epilepsy on DNA methylation. **Table S7.** Two-sample MR analysis of the effect of febrile and vaccine-related seizures on DNA methylation. **Table S8.** Top 50 associations with rs10258194 in MR_Base.

Abbreviations

ALSPAC: Avon longitudinal study of parents and children; ARIES: Accessible resource for integrative epigenomic studies; BDNF: Brain-derived neurotrophic factor; CpG: Cytosine-guanine dinucleotide; EWAS: Epigenome-wide association study; FDR: False discovery rate; GWAS: Genome-wide association study; IQR: Interquartile range; MMR: Mumps measles rubella; mQTL: Methylation quantitative trait loci; MR: Mendelian randomization; PheWAS: Phenome-wide association study; SM: Supplementary material; SNP: Single-nucleotide polymorphism; SV: Surrogate variable

Acknowledgements

We are extremely grateful to all the families who took part in the ALSPAC study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses.

The Generation R Study is conducted by the Erasmus Medical Center in close collaboration with the School of Law and Faculty of Social Sciences of the Erasmus University Rotterdam; the Municipal Health Service Rotterdam area, Rotterdam; the Rotterdam Homecare Foundation, Rotterdam; and the Stichting Trombosedienst & Artsenlaboratorium Rijnmond (STAR-MDC), Rotterdam. We gratefully acknowledge the contribution of children and parents, general practitioners, hospitals, midwives, and pharmacies in Rotterdam. The generation and management of the Illumina 450 K methylation array data (EWAS data) for the Generation R Study was executed by the Human Genotyping Facility of the Genetic Laboratory of the Department of Internal Medicine, Erasmus MC, the Netherlands. We thank Mr. Michael Verbiest, Ms. Mila Jhamai, Ms. Sarah Higgins, Mr. Marijn Verkerk and Dr. Lisette Stolk for their help in creating the EWAS database. We thank Dr. A. Teumer for his work on the quality control and normalization scripts.

Authors' contributions

DC designed the study, conducted the MR analysis, and revised the draft substantially; CH analyzed the ALSPAC data and wrote the initial draft; RHM analyzed the Generation R data; JFF acquired the Generation R data and assisted with data interpretation; CAC contributed substantially to the interpretation of the results; CLR, acquired the ALSPAC data and assisted with data interpretation; EW assisted with the design of the study, conducted some of the analyses, and contributed substantially to the interpretation of results. All authors read and contributed to the preparation of the final manuscript. All authors read and approved the final manuscript.

Funding

The UK Medical Research Council and Wellcome (Grant ref 102215/2/13/2) and the University of Bristol provide core support for ALSPAC. This publication is the work of the authors, and Doretta Caramaschi will serve as guarantors for the contents of this paper. A comprehensive list of grants funding is available on the ALSPAC website (<http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf>). GWAS data was generated by Sample Logistics and Genotyping Facilities at Wellcome Sanger Institute and LabCorp (Laboratory Corporation of America) using support from 23andMe. Methylation data in the ALSPAC cohort were generated as part of the UK BBSRC funded (BB/I025751/1 and BB/I025263/1) Accessible Resource for Integrated Epigenomic Studies (ARIES, <http://www.ariesepigenomics.org.uk>). D.C., C.R., and E.W. are funded by UK Medical Research Council (grant numbers: MC_UU_00011/1 and MC_UU_00011/5). C.H. is supported by a 4-year studentship fund from the Wellcome Trust

Molecular, Genetic and Lifecourse Epidemiology Ph.D. program at the University of Bristol (108902/B/15/Z).

The general design of the Generation R Study is made possible by financial support from the Erasmus Medical Center, Rotterdam, the Erasmus University Rotterdam, the Netherlands Organization for Health Research and Development and the Ministry of Health, Welfare and Sport. The EWAS data was funded by a grant from the Netherlands Genomics Initiative (NGI)/Netherlands Organisation for Scientific Research (NWO) Netherlands Consortium for Healthy Aging (NCHA; project nr. 050-060-810), by funds from the Genetic Laboratory of the Department of Internal Medicine, Erasmus MC, and by a grant from the National Institute of Child and Human Development (R01HD068437). J.F.F. has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 633595 (DynaHEALTH) and from the European Joint Programming Initiative "A Healthy Diet for a Healthy Life" (JPI HDHL, NutriPROGRAM project, ZonMw the Netherlands no.529051022). C.C. has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 707404. This project received funding from the European Union's Horizon 2020 research and innovation program (733206, LIFECYCLE).

Availability of data and materials

The participants' data used in this study are not publicly available due to privacy restrictions. However, the data may be available upon request following the relevant procedures for ALSPAC (<http://www.bristol.ac.uk/alspac/researchers/access/>) and Generation R (<https://generationr.nl/researchers/collaboration/>).

Ethics approval and consent to participate

Written informed consent has been obtained for all ALSPAC participants. Ethical approval for the ALSPAC study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees. Consent for biological samples has been collected in accordance with the Human Tissue Act (2004). Informed consent for the use of data collected via questionnaires and clinics was obtained from participants following the recommendations of the ALSPAC Ethics and Law Committee at the time. For the Generation R study, the study protocol was approved by the Medical Ethical Committee of the Erasmus Medical Centre, Rotterdam. Written informed consent was obtained for all participants.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Bristol Medical School, Population Health Sciences, University of Bristol, 5 Tyndall Avenue, Bristol BS8 1UD, UK. ²Medical Research Council Integrative Epidemiology Unit, University of Bristol, Oakfield House, Oakfield Grove, Bristol BS8 2BN, UK. ³Institute of Education and Child Studies, Leiden University, Pieter de la Court building, Wassenaarseweg 52, 2333, AK, Leiden, The Netherlands. ⁴Generation R Study Group, Erasmus MC, University Medical Center Rotterdam, Doctor Molewaterplein 40, 3015, GD, Rotterdam, The Netherlands. ⁵Department of Child and Adolescent Psychiatry/Psychology, Erasmus MC, University Medical Center Rotterdam, Doctor Molewaterplein 40, 3015, GD, Rotterdam, The Netherlands. ⁶Department of Pediatrics, Erasmus MC, University Medical Center Rotterdam, Doctor Molewaterplein 40, 3015, GD, Rotterdam, The Netherlands. ⁷Department of Epidemiology, Erasmus MC, University Medical Center Rotterdam, Doctor Molewaterplein 40, 3015, GD, Rotterdam, The Netherlands. ⁸Department of Psychology, University of Bath, Bath BA2 7AY, UK.

Received: 16 September 2019 Accepted: 2 December 2019

Published online: 08 January 2020

References

- Hauser WA. The prevalence and incidence of convulsive disorders in children. *Epilepsia*. 1994;35(Suppl 2):S1–6.
- Camfield P, Camfield C. Incidence, prevalence and aetiology of seizures and epilepsy in children. *Epileptic Disord*. 2015;17(2):117–23.

3. Scott RC, Tuchman R. Epilepsy and autism spectrum disorders: relatively related. *Neurology*. 2016;87(2):130–1.
4. Gillberg C, Lundstrom S, Fernell E, Nilsson G, Neville B. Febrile Seizures and epilepsy: association with autism and other neurodevelopmental disorders in the child and adolescent twin study in Sweden. *Pediatr Neurol*. 2017;74:80–6 e2.
5. Ku YC, Muo CH, Ku CS, Chen CH, Lee WY, Shen EY, et al. Risk of subsequent attention deficit-hyperactivity disorder in children with febrile seizures. *Arch Dis Child*. 2014;99(4):322–6.
6. Reilly C, Atkinson P, Das KB, Chin RF, Aylett SE, Burch V, et al. Neurobehavioral comorbidities in children with active epilepsy: a population-based study. *Pediatrics*. 2014;133(6):e1586–93.
7. Berg AT, Langfitt JT, Testa FM, Levy SR, DiMario F, Westerveld M, et al. Global cognitive function in children with epilepsy: a community-based study. *Epilepsia*. 2008;49(4):608–14.
8. Wo SW, Ong LC, Low WY, Lai PSM. The impact of epilepsy on academic achievement in children with normal intelligence and without major comorbidities: a systematic review. *Epilepsy Res*. 2017;136:35–45.
9. Abou-Khalil B, Auce P, Avbersek A, Bahlo M, Balding DJ, Bast T, et al. Genome-wide mega-analysis identifies 16 loci and highlights diverse biological mechanisms in the common epilepsies. *Nature Communications*. 2018;9(1):5269.
10. International League Against Epilepsy Consortium on Complex Epilepsies. Genetic determinants of common epilepsies: a meta-analysis of genome-wide association studies. *Lancet Neurol*. 2014;13(9):893–903.
11. Hauser RM, Henshall DC, Lubin FD. The epigenetics of epilepsy and its progression. *Neuroscientist*. 2018;24(2):186–200.
12. Scheffer IE, Berkovic S, Capovilla G, Connolly MB, French J, Guilhoto L, et al. ILAE classification of the epilepsies: position paper of the ILAE Commission for Classification and Terminology. *Epilepsia*. 2017;58(4):512–21.
13. Engel J Jr. Mesial temporal lobe epilepsy: what have we learned? *Neuroscientist*. 2001;7(4):340–52.
14. Long HY, Feng L, Kang J, Luo ZH, Xiao WB, Long LL, et al. Blood DNA methylation pattern is altered in mesial temporal lobe epilepsy. *Sci Rep*. 2017;7:43810.
15. Pathak S, Miller J, Morris EC, Stewart WCL, Greenberg DA. DNA methylation of the BRD2 promoter is associated with juvenile myoclonic epilepsy in Caucasians. *Epilepsia*. 2018;59(5):1011–9.
16. Xiao W, Cao Y, Long H, Luo Z, Li S, Deng N, et al. Genome-wide DNA methylation patterns analysis of noncoding RNAs in temporal lobe epilepsy patients. *Mol Neurobiol*. 2018;55(1):793–803.
17. Miller-Delaney SF, Bryan K, Das S, McKiernan RC, Bray IM, Reynolds JP, et al. Differential DNA methylation profiles of coding and non-coding genes define hippocampal sclerosis in human temporal lobe epilepsy. *Brain*. 2015; 138(Pt 3):616–31.
18. Kobow K, Kaspi A, Harikrishnan KN, Kiese K, Ziemann M, Khurana I, et al. Deep sequencing reveals increased DNA methylation in chronic rat epilepsy. *Acta Neuropathol*. 2013;126(5):741–56.
19. McDade TW, Ryan CP, Jones MJ, Hoke MK, Borja J, Miller GE, et al. Genome-wide analysis of DNA methylation in relation to socioeconomic status during development and early adulthood. *Am J Phys Anthropol*. 2019;169(1):3–11.
20. Chin RF, Neville BG, Peckham C, Wade A, Bedford H, Scott RC, et al. Socioeconomic deprivation independent of ethnicity increases status epilepticus risk. *Epilepsia*. 2009;50(5):1022–9.
21. Dingledine R, Coulter DA, Fritsch B, Gorter JA, Lelutiu N, McNamara J, et al. Transcriptional profile of hippocampal dentate granule cells in four rat epilepsy models. *Sci Data*. 2017;4:170061.
22. Davey Smith G, Hemani G. Mendelian randomization: genetic anchors for causal inference in epidemiological studies. *Hum Mol Genet*. 2014;23(R1):R89–98.
23. Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J, et al. Cohort Profile: The 'Children of the 90s'—the index offspring of the Avon Longitudinal Study of Parents and Children. *International Journal of Epidemiology*. 2013;42(1):111–27.
24. Fraser A, Macdonald-Wallis C, Tilling K, Boyd A, Golding J, Davey Smith G, et al. Cohort Profile: the Avon longitudinal study of parents and children: ALSPAC mothers cohort. *Int J Epidemiol*. 2013;42(1):97–110.
25. Brodie MJ, Zuberi SM, Scheffer IE, Fisher RS. The 2017 ILAE classification of seizure types and the epilepsies: what do people with epilepsy and their caregivers need to know? *Epileptic Disord*. 2018;20(2):77–87.
26. Relton CL, Gaunt T, McArdle W, Ho K, Duggirala A, Shihab H, et al. Data resource profile: accessible resource for integrated epigenomic studies (ARIES). *Int J Epidemiol*. 2015;44(4):1181–90.
27. Min JL, Hemani G, Davey Smith G, Relton C, Suderman M. Meffil: efficient normalization and analysis of very large DNA methylation datasets. *Bioinformatics*. 2018;34(23):3983–9.
28. Fortin JP, Labbe A, Lemire M, Zanke BW, Hudson TJ, Fertig EJ, et al. Functional normalization of 450 k methylation array data improves replication in large cancer studies. *Genome Biol*. 2014;15(12):503.
29. Marini C, King MA, Archer JS, Newton MR, Berkovic SF. Idiopathic generalised epilepsy of adult onset: clinical syndromes and genetics. *J Neurol Neurosurg Psychiatry*. 2003;74(2):192–6.
30. Barfield RT, Kilaru V, Smith AK, Conneely KN. CpGassoc: an R function for analysis of DNA methylation microarray data. *Bioinformatics*. 2012;28(9): 1280–1.
31. Teschendorff AE, Zhuang J, Widschwendter M. Independent surrogate variable analysis to deconvolve confounding factors in large-scale microarray profiling studies. *Bioinformatics*. 2011;27(11):1496–505.
32. Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH, et al. DNA methylation arrays as surrogate measures of cell mixture distribution. *BMC Bioinformatics*. 2012;13:16.
33. Bakulski KM, Feinberg JL, Andrews SV, Yang J, Brown S, McKenney SL, et al. DNA methylation of cord blood cell types: Applications for mixed cell birth studies. *Epigenetics*. 2016;11(5):354–62.
34. Kooijman MN, Kruithof CJ, van Duijn CM, Duijts L, Franco OH, van Ijzendoorn MH, et al. The Generation R Study: design and cohort update 2017. *Eur J Epidemiol*. 2016;31(12):1243–64.
35. Lehne B, Drong AW, Loh M, Zhang W, Scott WR, Tan ST, et al. A coherent approach for analysis of the Illumina HumanMethylation450 BeadChip improves data quality and performance in epigenome-wide association studies. *Genome Biol*. 2015;16:37.
36. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. 2010;26(17):2190–1.
37. Hemani G, Zheng J, Elsworth B, Wade KH, Haberland V, Baird D, et al. The MR-Base platform supports systematic causal inference across the human phenotype. *Elife*. 2018;7.
38. Gaunt TR, Shihab HA, Hemani G, Min JL, Woodward G, Lyttleton O, et al. Systematic identification of genetic influences on methylation across the human life course. *Genome Biol*. 2016;17:61.
39. Feenstra B, Pasternak B, Geller F, Carstensen L, Wang T, Huang F, et al. Common variants associated with general and MMR vaccine-related febrile seizures. *Nat Genet*. 2014;46(12):1274–82.
40. Sha'ari HM, Haerian BS, Baum L, Tan HJ, Rafia MH, Kwan P, et al. Association of BDNF polymorphisms with the risk of epilepsy: a multicenter study. *Mol Neurobiol*. 2016;53(5):2869–77.
41. Martinez-Levy GA, Rocha L, Lubin FD, Alonso-Vanegas MA, Nani A, Buentello-Garcia RM, et al. Increased expression of BDNF transcript with exon VI in hippocampi of patients with pharmaco-resistant temporal lobe epilepsy. *Neuroscience*. 2016;314:12–21.
42. Hannon E, Lunnon K, Schalkwyk L, Mill J. Interindividual methylomic variation across blood, cortex, and cerebellum: implications for epigenetic studies of neurological and neuropsychiatric phenotypes. *Epigenetics*. 2015; 10(11):1024–32.
43. Edgar RD, Jones MJ, Meaney MJ, Turecki G, Kobor MS. BECon: a tool for interpreting DNA methylation findings from blood in the context of brain. *Transl Psychiatry*. 2017;7(8):e1187.
44. Braun PR, Han S, Hing B, Nagahama Y, Gaul LN, Heinzman JT, et al. Genome-wide DNA methylation comparison between live human brain and peripheral tissues within individuals. *Transl Psychiatry*. 2019;9(1):47.
45. Walsh S, Donnan J, Fortin Y, Sikora L, Morrissey A, Collins K, et al. A systematic review of the risks factors associated with the onset and natural progression of epilepsy. *Neurotoxicology*. 2017;61:64–77.
46. Okbay A, Beauchamp JP, Fontana MA, Lee JJ, Pers TH, Rietveld CA, et al. Genome-wide association study identifies 74 loci associated with educational attainment. *Nature*. 2016;533(7604):539–42.
47. Benyamin B, Pourcain B, Davis OS, Davies G, Hansell NK, Brion MJ, et al. Childhood intelligence is heritable, highly polygenic and associated with FBNP1L. *Mol Psychiatry*. 2014;19(2):253–8.
48. Cross-Disorder Group of the Psychiatric Genomics C, Lee SH, Ripke S, Neale BM, Faraone SV, Purcell SM, et al. Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. *Nat Genet*. 2013; 45(9):984–94.
49. Jankowsky JL, Patterson PH. The role of cytokines and growth factors in seizures and their sequelae. *Prog Neurobiol*. 2001;63(2):125–49.

50. Kandratavicius L, Monteiro MR, Assirati JA Jr, Carlotti CG Jr, Hallak JE, Leite JP. Neurotrophins in mesial temporal lobe epilepsy with and without psychiatric comorbidities. *J Neuropathol Exp Neurol*. 2013;72(11):1029–42.
51. Pruunsild P, Kazantseva A, Aid T, Palm K, Timmusk T. Dissecting the human BDNF locus: bidirectional transcription, complex splicing, and multiple promoters. *Genomics*. 2007;90(3):397–406.
52. Ozdemir O, Egemen E, Ugur Iseri SA, Sezerman OU, Bebek N, Baykan B, et al. Identification of epilepsy related pathways using genome-wide DNA methylation measures: a trio-based approach. *PLoS One*. 2019;14(2): e0211917.
53. Anney R, Klei L, Pinto D, Regan R, Conroy J, Magalhaes TR, et al. A genome-wide scan for common alleles affecting risk for autism. *Hum Mol Genet*. 2010;19(20):4072–82.
54. Lee JJ, Wedow R, Okbay A, Kong E, Maghzian O, Zacher M, et al. Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nat Genet*. 2018;50(8): 1112–21.
55. Jansen S, van der Werf IM, Innes AM, Afejar A, Agrawal PB, Anderson LJ, et al. De novo variants in FBXO11 cause a syndromic form of intellectual disability with behavioral problems and dysmorphisms. *Eur J Hum Genet*. 2019.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

